



Recombinant Mouse TFF3 (C-6His)

Catalog #	EPT168
Expression Host	Human Cells
DESCRIPTION	Recombinant Mouse Trefoil Factor 3 is produced by our Mammalian expression system and the target gene encoding Ala23-Phe81 is expressed with a 6His tag at the C-terminus.
Accession	Q62395
Synonyms	Trefoil factor 3; Intestinal trefoil factor; mITF; Tff3; Itf
Mol Mass	7.3 KDa
AP Mol Mass	11 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.





It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

BACKGROUND

Trefoil factors (TFF) are secretory products of mucin producing cells. They play a key role in the maintenance of the surface integrity of oral mucosa and enhance healing of the gastrointestinal mucosa by a process called restitution. TFF comprises the gastric peptides (TFF1), spasmolytic peptide (TFF2), and the intestinal trefoil factor (TFF3). They have an important and necessary role in epithelial restitution within the





gastrointestinal tract. Members of the trefoil family are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfide bonds. They are stable secretory proteins expressed in gastrointestinal mucosa. Trefoil Factor 3(TFF3) is involved in the maintenance and repair of the intestinal mucosa. TFF3 promotes the mobility of epithelial cells in healing processes (motogen).

SDS-PAGE

